

Pain Management Coordination

PROVIDER TEMPLATES FOR PATIENTS' TAPS SCREENING

July 2022

These templates are meant to notify Surgeons and Surgical Care Teams of patients with a positive screen and recommendations specific to the area of risk. For patients with positive screens for multiple categories, both categories' resources can be cohesively combined and sent in one notification message. Prescription and non-prescription opioids have the same templates. Prescription and non-prescription stimulants have the same templates.

CONTENTS

SCORING CATEGORY CRITERIA	1
SURGEON MESSAGE TEMPLATES	2
General Positive Screen	2
Tobacco	2
Alcohol	3
Sedatives	4
Cannabis	5
Stimulants	6
Opioids	7
Other Substances	8
ADVANCED PRACTICE PROVIDER MESSAGE TEMPLATES	9
General Positive Screen	9
Tobacco	10
Alcohol	11
Sedatives	12
Cannabis	13
Stimulants	14
Opioids	15
Other Substances	16

Click on the title above to jump to the page in this document.



SCORING CATEGORY CRITERIA

Scoring Category:		Positive Screen Criteria:	Total Possible Score:
SC0	Tobacco	2	3
SC1	Alcohol	2	4
SC2	Cannabis	2	3
SC3	Stimulant	1	3
SC4	Heroin	1	3
SC5	Opioid	1	3
SC6	Sedative	1	3
SC7	Rx Stimulant	1	3
SC8	Other Substance	Self-report any substance	NA

Scoring is based on the TAPS 2 screening assessment: [PubMed Central Image Viewer](#).

Online screening tool: <https://nida.nih.gov/taps2/>

MESSAGE TEMPLATES

General Positive Screen

To:	Surgeon and/or Surgical Care Team
Subject Line:	Please review: Your Surgical Patient's TAPS screening
<p><u>Surgeon name</u> .</p> <p>Your patient (<u>MRN</u> ; DOS <u>[date of surgery]</u>) is at risk for adverse outcomes related to <u>[substance]</u> and may meet criteria for a <u>category</u> . Hearing from their surgeon can be helpful to manage this risk.</p> <p><u>[Insert all screened positive category content]</u></p> <p>Your patient completed the Tobacco, Alcohol, Prescription medications and other Substance Tools (TAPS) screener on <u>date</u></p>	

Tobacco

[insert Tobacco Use Disorder into category]

Suggested Action

1. CONFIRM diagnosis through clinical interview and review of tobacco use

2. EXPRESS CONCERN using empathy. Discuss any [perioperative risks](#) and determine if patient desires tobacco cessation resources

3. CREATE A PERIOPERATIVE PLAN proactively

- Offer and provide tobacco cessation [resources](#) to patient, if patient agrees
- Refer to MHealthy Tobacco Cessation Services, if patient agrees
- Communicate plan with the patient, patient's primary care provider, surgical team, anesthesia team

4. EDUCATE/ADVISE patient on:

- Educate patient on holding tobacco products minimally, 2 weeks prior to surgery

Additional Resources:

- Tobacco Cessation [resources](#)
- [Nicotine Replacement Therapy](#)

Alcohol

[insert Alcohol Use Disorder into category]

Suggested Action

1. CONFIRM diagnosis through clinical interview, review alcohol use, and assess for other co-occurring problems that may be impacting alcohol use (e.g. pain, anxiety, depression)

2. EXPRESS CONCERN using empathy. Recommend reducing alcohol consumption and discuss associated increased risk of [perioperative complications](#)

3. CREATE A PERIOPERATIVE PLAN proactively

- Outpatient: Consult Addiction Treatment Services for possible tapering plan and/or withdrawal management
 - In basket to request appt. Can be done through: P rub Psy Adult Admn Sup
- Inpatient: Consult Addiction Consult Team for possible tapering plan and/or withdrawal management
 - Per [UM guidelines](#), upon transfer to floor, all adult inpatients for risk of alcohol withdrawal will be screened by Nursing using AUDIT-C
- Recommend participation in [Alcohol Management Program](#)
- Communicate plan with the patient, patient's primary care provider, surgical team, anesthesia team, and other disciplines (e.g. pain service, addiction service, psych consult team)

4. EDUCATE/ADVISE patient on:

- [Benefits of reduced alcohol](#) consumption
- Risks associated with combining [alcohol with other drugs](#) (especially opioids, benzodiazepines)
- Against alcohol use and driving
- Following up with Primary Care Physician for counseling and/or support group to complement [FDA medication treatment](#), when warranted

Additional Resources

- [AAFP: Outpatient Management of Alcohol Withdrawal Syndrome](#)
- [Michigan Opioid Collaborative consult services](#)

Sedatives

[insert Sedative Use Disorder into category]

Suggested Action

1. CONFIRM diagnosis through clinical interview, review prescription and non-prescription sedative use, and assess for other co-occurring problems (e.g. pain, anxiety, depression)

2. EXPRESS CONCERN using empathy. Recommend cessation of non-prescribed sedative use and discuss associated increased risk of perioperative complications

3. CREATE A PERIOPERATIVE PLAN proactively

- Outpatient: Consult Addiction Treatment Services for possible tapering plan and/or withdrawal management
 - In basket to request appt. can be done through: P rub Psy Adult Admn Sup
- Inpatient: Consult Addiction Consult Team for possible tapering plan and/or withdrawal management
- Order a Point of Care Urine Drug Screen during the preoperative clinic visit and on the Day of Surgery, if patient endorses recent illicit sedative use
- Communicate plan with the patient, patient's usual prescriber, surgical team, anesthesia team, and other disciplines (e.g. pain service, addiction service, psych consult team)

4. EDUCATE/ADVISE patient on:

- Danger of abrupt discontinuation of benzodiazepines
- Increased risk when mixing sedatives with opioids and/or alcohol
- Against sedative use and driving
- Following up with Primary Care Physician to discuss tapering the use of prescribed sedative, possible switching sedative, and referral to counseling or treatment program for substance use disorder and any co-occurring problems, such as depression, that may be driving sedative use.

Additional Resources

- [SAMHSA: Behavioral Health Treatment Services Locator](#)
- [SAMHSA: Detoxification and Substance Abuse Treatment](#)
- [Michigan Opioid Collaborative consult services](#)

Cannabis

[insert Cannabis Use Disorder into category]

Suggested Action

1. CONFIRM diagnosis through clinical interview, and assess for other co-occurring problems (e.g. pain, anxiety, depression).

2. EXPRESS CONCERN using empathy. Recommend reduction and/or cessation of cannabis and discuss associated increased risk of perioperative complications.

3. CREATE A PERIOPERATIVE PLAN proactively:

- Review any prescribed medications to identify those that may increase the risk of sedation and impairment (opioids, benzodiazepines).

4. EDUCATE/ADVISE patient on:

- Educate patient on holding cannabis minimally 1 week prior to surgery
- Risks associated with combining cannabis with alcohol or other drugs.
- Against cannabis use and driving.
- Following up with a Primary Care Physician for a referral to counseling or a treatment program for SUD and any co-occurring problems, such as depression, that may be impacting cannabis use.

Stimulants

[insert Stimulant Use Disorder into category]

Suggested Action

1. CONFIRM diagnosis through clinical interview, review prescription and non-prescription stimulant use and assess for other co-occurring problems (e.g. pain, anxiety, depression)

2. EXPRESS CONCERN using empathy and discuss associated increased risk of perioperative complications

3. CREATE A PERIOPERATIVE PLAN proactively

- Outpatient: Consult Addiction Treatment Services for possible tapering plan and/or withdrawal management
 - In basket to request appt. can be done through: P rub Psy Adult Admn Sup
- Inpatient: Consult Addiction Consult Team for possible tapering plan and/or withdrawal management
- Order a Point of Care Urine Drug Screen during the patient's preoperative clinic visit and on the Day of Surgery, if patient admits to recent illicit stimulant use
- Communicate plan with the patient, patient's outpatient prescriber, surgical team, anesthesia team, and other disciplines (e.g. pain service, addiction service, psych consult team)

4. EDUCATE/ADVISE patient on:

- Educate patient on reducing or stopping stimulant use before surgery
- On risks associated with injecting stimulants with heroin or otherwise combining stimulants with alcohol or other drugs.
- Against stimulant use and driving
- Following up with a Primary Care Physician for behavioral health counseling referral coupled with pharmacotherapy for most effective treatment.
- [Harm Reduction](#)

Additional Resources

- [SAMHSA: Behavioral Health Treatment Services Locator](#)
- [NIDA: Principles of Drug Addiction Treatment: A Research-Based Guide \(3rd ed.\) - Behavioral Therapies](#)
- [Michigan Opioid Collaborative consult services](#)

Opioids

[insert Opioid Use Disorder into category]

Suggested Action

1. CONFIRM diagnosis through clinical interview, review prescription and non-prescription opioid use and assess for other co-occurring problems (e.g. pain, anxiety, depression)

2. EXPRESS CONCERN using empathy and discuss the [perioperative period risks](#)

3. CREATE A PERIOPERATIVE PLAN proactively

- Consider inpatient postoperative admission for pain management which is challenging in this population
- Inpatient: Consult Addiction Consult Team (ACT) for possible treatment
- Outpatient: Consult Addiction Treatment Services for possible treatment
 - In basket to request appt. can be done through: P rub Psy Adult Admn Sup
- Consult:
 - In-house preoperative anesthesia to determine if an Acute Pain Service referral is warranted if patient takes >100mme/day
 - Acute Pain Services (APS) to address complex pain management concerns
- Refer to Michigan Medicine's MOUD protocol and policies if the patient is currently taking any Medications for Opioid Use Disorder (buprenorphine, suboxone, methadone, naltrexone)
- Consider prescribing [naloxone](#)
- Use evidence-based postoperative [opioid prescribing recommendations](#) and [non-opioid recommendations](#)
- Communicate plan with the patient, patient's outpatient prescriber, surgical team, anesthesia team, and other disciplines (e.g. pain service, addiction service, psych consult team)

4. EDUCATE/ADVISE patient on:

- Postoperative pain expectations
- Postoperative [over the counter pain management](#)
- Legislatively mandated [opioid pain management](#) education
- [Overdose prevention](#)
- [Harm Reduction](#)
- Following up with Primary Care Physician for Medications for Opioid Use Disorder (MOUD) and behavioral health counseling referral

Additional Resources

- [SAMHSA Behavioral Health Treatment Services Locator](#)
- [Michigan Opioid Collaborative consult services](#)

Other Substances

[insert Problem Substance Use into category]

Suggested Action

1. CONFIRM use of illicit/recreational drugs through clinical interview. Review illicit drug use, prescribed medications, and assess for other co-occurring problems (e.g. pain, anxiety, and depression)

2. EXPRESS CONCERN using empathy. Recommend cessation, and discuss associated increased risk of perioperative complications

3. CREATE A PERIOPERATIVE PLAN proactively

- If patient admits to recent use of Molly or LSD, order a Point Of Care Urine Drug Screen during the preoperative clinic visit and on the Date Of Surgery
- Discuss with in-house Preoperative Anesthesia as needed
- Communicate with the patient, surgical team, anesthesia team, and other disciplines (e.g. pain service, addiction service, psych consult team)

4. EDUCATE/ADVISE patient on:

- Against illicit/recreational drug use and driving
- Mixing illicit/recreational drugs with their postoperative opioid prescription (if applicable)
- Following up with Primary Care Physician for behavioral health/counseling referral