

REFERENCES

1. Maughan B. Drug and alcohol dependence, Elsevier. 2016; 168:328-34.
2. Larach D. Ann Surg. 2020;271(2):290-295.
3. Chua, K-P. AAP, 2021;148(3), e2021051539.
4. Chua, K-P. AJPM 2021;61(2), 165–173.
5. Nalliah R. JAMA Network Open. 2020;3(3):e200901.
6. Howard R. JAMA Surg. 2018;153(3):285-287.
7. Moore P. JADA. 2018;149(4):256-65.
8. American Dental Association. (2022.) Oral Analgesics for Acute Dental Pain.
9. Schroeder A. JAMA Internal Medicine. 2019;179(2):145-52.
10. Harbaugh C. JAMA Network. 2018;320(5):504-6.
11. Johnston L. Ann Arbor, MI: The University of Michigan; 2016.

Scan for OPEN Healthcare
Professional resources



OPEN is partially funded by the Michigan
Department of Health and Human Services
<https://doi.org/10.56137/OPEN.000016>
Updated 3.24

IMPROVE OPIOID PRESCRIBING PRACTICES AFTER DENTAL SURGERY



4 EVIDENCE- BASED REASONS

MICHIGAN-OPEN.ORG

OPEN

Prevention. Treatment. Recovery.

EVIDENCE SHOWS:

In outpatient dental surgery,
54% of opioids go unused¹

Reducing opioid prescribing improves the safety for patients, families, and communities

- From 2010 to 2016, the proportion of initial opioid prescriptions in dental care increased by 68%²
- In young adults, dentists are the leading prescribers of opioids³
- Overdose risk among family members increased when opioid prescriptions were dispensed to patients for a dental surgery⁴

NO CORRELATION

between patient satisfaction scores and amount of opioid prescribed⁵

Prescribing more opioids does not improve patient satisfaction

- Patients who used opioids after tooth extraction reported significantly higher levels of pain compared to those who did not use opioids, but no difference in satisfaction was observed⁵
- Patients who were prescribed fewer opioids reported using fewer opioids with no change in pain scores⁶

Acetaminophen and ibuprofen are **more effective** than opioids in managing pain^{7,8}



Using OTC medications to manage pain can decrease the need for opioids after dental surgery

- A combination of ibuprofen and acetaminophen is more effective than opioids for dental pain control and carries less risk for adverse events^{7,8}
- The ADA recommends NSAIDs as a first line therapy⁸

6.9% Of adolescents & young adults who receive opioids after dental surgery developed new persistent opioid use⁹

New persistent opioid use is one of the most common surgical complications

- In adolescents and young adults who received dental opioids, 5.8% had an opioid misuse or overdose related encounter⁹
- Filling an opioid prescription after wisdom tooth removal resulted in more than three times the risk of developing new persistent opioid use¹⁰
- Most adolescents believe that prescription opioids are safer than other substances of misuse¹¹

OPEN
Prescribing
Recommendations

